

Solution of 2016 Midterm (Math 2033)

① We have $A \subseteq [3, 6]$ and $x \in [5, \sqrt{80})$. So $3 \leq y \leq 6$ and $5 \leq x < \sqrt{80}$.
Then $9 \leq y^2 \leq 36$ and $9 \leq x^2 - 16 < 64$ ($\Leftrightarrow 3 \leq \sqrt{x^2 - 16} < 8$). So
 $12 \leq y^2 + \sqrt{x^2 - 16} < 44$. Hence B is bounded above by 44 and below by 12.

By supremum limit theorem, $\exists y_n \in A$ such that $\lim_{n \rightarrow \infty} y_n = 6$.

By infimum limit theorem, $\exists y'_n \in A$ such that $\lim_{n \rightarrow \infty} y'_n = 3$.


Next let $x_n = \sqrt{80} - \frac{1}{n}$ and $x'_n = 5 + \frac{1}{n\sqrt{2}}$, then $x_n, x'_n \in [5, \sqrt{80}) \setminus \mathbb{Q}$
and $\lim_{n \rightarrow \infty} x_n = \sqrt{80}$, $\lim_{n \rightarrow \infty} x'_n = 5$. Then $y_n^2 + \sqrt{x_n^2 - 16} \in B$ and $\lim_{n \rightarrow \infty} y_n^2 + \sqrt{x_n^2 - 16} = 44$.

Also $y_n'^2 + \sqrt{x_n'^2 - 16} \in B$ and $\lim_{n \rightarrow \infty} y_n'^2 + \sqrt{x_n'^2 - 16} = 12$. $\therefore \sup B = 44$ and $\inf B = 12$.

② Sketch: $x_1 = 35, x_2 = 3, x_3 = 15, x_4 = 6$

$$x = \frac{120}{5+x}$$

$$\Leftrightarrow x^2 + 5x - 120 = 0 \Leftrightarrow x = \frac{-5 \pm \sqrt{25 + 480}}{2} = \frac{-5 \pm \sqrt{505}}{2} \text{ (reject -)}$$



Solution Let $I_n = [x_{2n}, x_{2n-1}]$ for $n=1, 2, 3, \dots$

Claim: $I_n \supseteq I_{n+1}$ ($\Leftrightarrow x_{2n} \leq x_{2n+2} \leq x_{2n+1} \leq x_{2n-1}$) for $n=1, 2, 3, \dots$

Case $n=1$, $x_2 = 3 \leq x_4 = 6 \leq x_3 = 15 \leq x_1 = 35$

Suppose Case n is true. Then $x_{2n} \leq x_{2n+2} \leq x_{2n+1} \leq x_{2n-1}$. We have

$$5 + x_{2n} \leq 5 + x_{2n+2} \leq 5 + x_{2n+1} \leq 5 + x_{2n-1} \Rightarrow \frac{1}{5+x_{2n}} \geq \frac{1}{5+x_{2n+2}} \geq \frac{1}{5+x_{2n+1}} \geq \frac{1}{5+x_{2n-1}}$$

$$\Rightarrow x_{2n+1} = \frac{120}{5+x_{2n}} \geq x_{2n+3} = \frac{120}{5+x_{2n+2}} \geq x_{2n+2} = \frac{120}{5+x_{2n+1}} \geq x_{2n} = \frac{120}{5+x_{2n-1}}$$

$$\Rightarrow 5 + x_{2n+1} \geq 5 + x_{2n+3} \geq 5 + x_{2n+2} \geq 5 + x_{2n} \Rightarrow \frac{1}{5+x_{2n+1}} \leq \frac{1}{5+x_{2n+3}} \leq \frac{1}{5+x_{2n+2}} \leq \frac{1}{5+x_{2n}}$$

$$\Rightarrow x_{2n+2} = \frac{120}{5+x_{2n+1}} \leq x_{2n+4} = \frac{120}{5+x_{2n+3}} \leq x_{2n+3} = \frac{120}{5+x_{2n+2}} \leq x_{2n+1} = \frac{120}{5+x_{2n}}$$

This completes M.I. So the claim is true.

By nested interval theorem, $\lim_{n \rightarrow \infty} x_{2n} = a$ and $\lim_{n \rightarrow \infty} x_{2n-1} = b$. Now $\lim_{n \rightarrow \infty} x_{2n+1} = \lim_{n \rightarrow \infty} \frac{120}{5+x_{2n}}$

$$\Rightarrow b = \frac{120}{5+a} \text{ and } \lim_{n \rightarrow \infty} x_{2n} = \lim_{n \rightarrow \infty} \frac{120}{5+x_{2n-1}} \Rightarrow a = \frac{120}{5+b}. \text{ We have } 5b + ab = 120 = 5a + ab.$$

This implies $a = b$. Then $a = \lim_{n \rightarrow \infty} x_{n+1} = \lim_{n \rightarrow \infty} \frac{120}{5+x_n} = \frac{120}{5+a} \Rightarrow a^2 + 5a - 120 = 0$

$$\Rightarrow a = \frac{-5 \pm \sqrt{25 + 480}}{2} = \frac{-5 \pm \sqrt{505}}{2}. \text{ Since } x_n \geq 3, a = \lim_{n \rightarrow \infty} x_n \text{ is } \frac{-5 + \sqrt{505}}{2}.$$

③ $S = \{r\sqrt{2} : r \in \mathbb{Q}\} = \bigcup_{r \in \mathbb{Q}} \underbrace{\{r\sqrt{2}\}}_{\substack{\text{1 element} \\ \Rightarrow \text{countable}}}$ is countable by countable union theorem.

Let $W = \{2^x \sin y - 2^y \cos x : x, y \in S\}$. Then $W = \bigcup_{(x,y) \in S \times S} \underbrace{\{2^x \sin y - 2^y \cos x\}}_{\substack{\text{1 element} \\ \Rightarrow \text{countable}}}$ is countable. Then $\underbrace{\mathbb{R}}_{\text{uncountable}} \setminus \underbrace{W}_{\text{countable}}$ is uncountable. So $\mathbb{R} \setminus W$ is an infinite set. Therefore, there exist infinitely many real numbers c such that $2^x \sin y - 2^y \cos x = c$ has no solution with $x, y \in S$.

④ Sketch $\frac{4n\sqrt{n}-3}{n\sqrt{n}+5} \rightarrow 4, \frac{6n^2-2n+3}{3n^2-1} \rightarrow 2$

$$\left\{ \begin{aligned} \left| \frac{4n\sqrt{n}-3}{n\sqrt{n}+5} - 4 \right| &= \frac{|-23|}{n\sqrt{n}+5} \leq \frac{23}{n\sqrt{n}} = \frac{23}{n^{3/2}} < \frac{\varepsilon}{2} \Leftrightarrow \left(\frac{46}{\varepsilon}\right)^{2/3} < n \\ \left| \frac{6n^2-2n+3}{3n^2-1} - 2 \right| &= \frac{|-n+5|}{3n^2-1} \leq \frac{n+5}{3n^2-n^2} \leq \frac{6n}{2n^2} = \frac{3}{n} < \frac{\varepsilon}{2} \Leftrightarrow \frac{6}{\varepsilon} < n \end{aligned} \right.$$

Solution For every $\varepsilon > 0$, by Archimedean principle, $\exists K \in \mathbb{N}$ such that $K > \max\left\{\left(\frac{46}{\varepsilon}\right)^{2/3}, \frac{6}{\varepsilon}\right\}$ (or take $K = \lceil \max\left\{\left(\frac{46}{\varepsilon}\right)^{2/3}, \frac{6}{\varepsilon}\right\} \rceil$). Then $n \geq K \Rightarrow \left| \frac{4n\sqrt{n}-3}{n\sqrt{n}+5} + \frac{6n^2-2n+3}{3n^2-1} - 6 \right| \leq \left| \frac{4n\sqrt{n}-3}{n\sqrt{n}+5} - 4 \right| + \left| \frac{6n^2-2n+3}{3n^2-1} - 2 \right| < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon.$